

(19)



Europäisch s Pat ntamt

Europ an Patent Office

Office européen des br v ts



(11)

**EP 1 099 444 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

16.05.2001 Bulletin 2001/20

(51) Int. Cl.<sup>7</sup>: **A61K 35/78**, C07J 7/00,

C07J 17/00

(21) Application number: **00309271.5**

(22) Date of filing: **20.10.2000**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **27.10.1999 GB 9925457**

(71) Applicant: **PHYTOPHARM PLC**

**Godmanchester, Cambridgeshire PE18 8HG  
(GB)**

(72) Inventors:

- **Horak, Roelof Marthinus**  
**0181 Pretoria (SA)**
- **Maharaj, Vinesh**  
**Heuwelord 0149 (SA)**
- **Hakkinen, John**  
**North Stonington, CT 06359 (US)**

(74) Representative:

**Abrams, Michael John et al**  
**Haseltine Lake & Co.,**  
**Imperial House,**  
**15-19 Kingsway**  
**London WC2B 6UD (GB)**

(54) **Gastric acid secretion**

(57) The present application relates to the reduction of gastric acid secretion in animals including humans; to methods of treating conditions or disorders associated with or exacerbated by gastric acid secretion; to pharmaceutical compositions useful in carrying out such methods; and to the use of certain extracts and compounds in the treatment of human and non-human animals, generally mammals.

**EP 1 099 444 A1**


## Description

[0001] This invention relates to the reduction of gastric acid secretion in animals including humans; to methods of treating conditions or disorders associated with or exacerbated by gastric acid secretion; to pharmaceutical compositions useful in carrying out such methods; and to the use of certain extracts and compounds in the treatment of human and non-human animals, generally mammals.

[0002] Excessive secretion of gastric acid can lead to or aggravate a number of disorders, for example oesophageal reflux disease, e.g. reflux oesophagitis, haemorrhage in and benign ulceration of the stomach and duodenum (including those complicating NSAID therapy). These conditions tend to be more problematical in obese patients and patients with hiatus hernia. The present invention is directed, in part, to the treatment of these conditions and of ancillary indications, e.g. to provide relief of reflux-like symptoms (e.g. heartburn) and/or ulcer-like symptoms (e.g. epigastric pain) associated with acid-related dyspepsia; for general dyspeptic symptoms; and for prophylaxis of acid aspiration.

[0003] Non-steroidal anti-inflammatory drugs (NSAIDs), e.g. aspirin (acetylsalicylic acid), are well-known and widely used for their anti-inflammatory and antipyretic properties. A major side effect is their tendency to damage the wall of the stomach; this adverse property is acid-dependent and is generic to the NSAIDs. The present invention is also concerned with means for mitigating these adverse effects.

[0004] It is known from International Patent Publication No. WO 98/46243 that extracts of certain plants of the genus *Trichocaulon* or *Hoodia* possess appetite suppressant properties. This document also discloses certain specific compounds which possess appetite suppressant activity. Among these is the compound 3-O-[- $\beta$ -D-thevetopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one; the structural formula of this compound is given as formula (1) in WO 98/46243. We have found that this compound is effective in reducing the secretion of gastric acid; accordingly, the compound finds application in the present invention. Derivatives of this compound are also effective in the present invention; such derivatives have the general formula



in which

R = alkyl;


R<sub>1</sub> = H, alkyl, tigloyl, benzoyl, or any other organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrate, or one or more 2,6-dideoxy carbohydrates, or glucose molecules, or combinations thereof;

in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-; and in which the dashed lines indicate the optional presence of a double bond in the C4-C5 or C5-C6 positions.

Thus, according to a first aspect of the present invention, there is provided a method of reducing gastric acid secretion in an animal, which comprises administering to said animal:

- (a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or
- (b) a compound of the formula



wherein:

R = an alkyl group containing from one to four carbon atoms;

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;

and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

[0005] Advantageously, when R<sub>1</sub> is an organic ester group it is tigloyl, benzoyl or anthraniloyl. In a preferred embodiment, R is an alkyl group having from one to four carbon atoms, R<sub>1</sub> is tigloyl or anthraniloyl; and a double bond is present in the C5-C6 position.


[0006] Advantageously, R<sub>2</sub> is trisaccharide group. The component sugars of said trisaccharide group are preferably 6-deoxy and/or 2,6-dideoxy hexoses. In some embodiments, the terminal hexose moiety is thevetosyl.

[0007] The compound may be 3-O-[-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one.

[0008] According to a second aspect of the present invention, there is provided a method of treating a disorder of the alimentary system in an animal, which comprises administering to the animal

(a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or

(b) a compound of the formula



wh rein:

R = an alkyl group containing from one to four carbon atoms;

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;


and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

[0009] This method can conveniently be used in the treatment of reflux oesophagitis. It is also expected to find application in the treatment of epigastric pain; dyspepsia; gastric ulceration; and acid aspiration.

[0010] According to a third aspect of the present invention, there is provided a method of protecting a mammalian gastro-intestinal tract from damage caused by a non-steroidal anti-inflammatory drug, which comprises administering an effective amount of:

(a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or

(b) a compound of the formula



wherein:

R = an alkyl group containing from one to four carbon atoms;

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;


and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

[0011] According to a fourth aspect, the present invention provides a method of protecting a mammalian gastro-intestinal tract from damage caused by a non-steroidal anti-inflammatory drug, which comprises administering an effective amount of 13-O-[-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a derivative thereof.

[0012] When a plant extract is used in the invention, it is presently preferred to use an extract from the plant *Hoodia gordonii* or *Hoodia currori*. Advantageously, the said extract comprises spray-dried sap of the plant *Hoodia gordonii*.

[0013] The methods in accordance with this invention are applicable to the treatment of humans.

[0014] According to a fifth aspect of the present invention, there is provided a pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a compound of the formula



wherein:

R = an alkyl group containing from one to four carbon atoms;

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;


in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;

and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

**[0015]** A preferred pharmaceutical composition of this type utilises the compound 3-O-[-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a derivative thereof. The pharmaceutical compositions of this invention are conveniently prepared in unit dosage form; they may also include a pharmaceutically acceptable diluent, excipient or carrier.

**[0016]** According to a sixth aspect of the present invention, there is provided method of treating a condition or disorder caused by or exacerbated by gastric acid secretion in an animal, which comprises administering to the animal a pharmaceutical composition as just defined. The animal may be a human.

**[0017]** According to a seventh aspect, the present invention provides the use of a compound of the formula



wherein:

R = an alkyl group containing from one to four carbon atoms;

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

$R_2 = H$ , or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

in which the acyl group  $R-(C=O)-$  group can be in the reduced form  $R-(C-OH)-$ ;

and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position,

in the manufacture of a medicament for the treatment of a disorder caused by or exacerbated by gastric acid secretion. For such use, the compound may be 3-O-[- $\beta$ -D-thevetopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one or a derivative thereof.

**[0018]** According to an eighth aspect, the present invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for the treatment of conditions and disorders caused by or exacerbated by gastric acid secretion. The condition or disorder for which the medicament is intended may be reflux oesophagitis; or gastric or duodenal ulceration; gastro-duodenal erosion; or epigastric pain.

**[0019]** According to a ninth aspect, the present invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for the treatment of conditions and disorders caused by or exacerbated by gastric acid secretion.

**[0020]** According to a tenth aspect, the present invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for the treatment of reflux oesophagitis.

**[0021]** According to an eleventh aspect, the present invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for preventing or reducing gastric damage associated with use of a non-steroidal anti-inflammatory drug.

**[0022]** According to a twelfth aspect, the present invention provides the use of 3-O-[- $\beta$ -D-thevetopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one or a derivative thereof in the manufacture of a medicament for preventing or reducing gastric damage associated with use of a non-steroidal anti-inflammatory drug.

**[0023]** The extracts of a plant of the genus *Hoodia* or *Trichocaulon* useful in the present invention may generally be prepared by extracting sap from the plant and then spray-drying the sap. Alternatively, solvent extraction procedures may be employed. In either case, fractionation of the initial extract, e.g. by column chromatography, may follow in order to generate an extract with enhanced activity.


**[0024]** The extract may be prepared from plant material such as the stems and roots of plants of the genus *Hoodia* or the genus *Trichocaulon*; these genera include succulent plants which grow in the arid regions of southern Africa. Advantageously, the plant extract is obtained from one of the following species: *Trichocaulon piliferum*; *Trichocaulon officinale*; *Hoodia currorii*; *Hoodia gordonii*; and *Hoodia lugardii*.

**[0025]** The plant material may be homogenised in the presence of a suitable solvent, e.g. a methanol/methylene chloride solvent, by means of a device such as a Waring blender. The extraction solution may then be separated from residual plant material by an appropriate separation procedure such as, for example, filtration or centrifugation. The solvent may, for example, be removed by means of a rotary evaporator, preferably in a water bath at a temperature of 60°C. The separated crude extract may then be further extracted with methylene chloride and water before being separated into a methylene chloride extract and a water extract. The methylene chloride extract may have the solvent removed by, for example, a rotary evaporator and the resultant extract may be further purified by way of a methanol/hexane extraction. The methanol/hexane extraction product may then be separated to yield a methanol extract and a hexane extract. The methanol extract may be evaporated to remove the solvent in order to yield a partially purified active extract.

**[0026]** The partially purified active extract may be dissolved in methanol, and may be further fractionated by column chromatography, employing silica gel as an adsorption medium and a chloroform/30% methanol mixture as eluent. A plurality of different fractions may be obtained, and each may be evaluated, by suitable bioassaying procedures, to determine their activity. High activity fractions may be further fractionated, e.g. by column chromatography using silica gel as an adsorption medium and a 9:1 chloroform:methanol eluent. This process may be repeated using silica gel as an adsorption medium and a 9:1 ethylacetate:hexane eluent.

**[0027]** Alternatively, the plant product may be compressed or macerated to extract sap therefrom, then filtered to remove unwanted solids and freeze-dried. The freeze-dried sap product may then be further purified, if desired, for example using chromatographic fractionation such as described above.

**[0028]** The derivatives of 3-O-[- $\beta$ -D-thevetopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one which find use in this invention are advantageously of the formula:



in which

R = alkyl;

R<sub>1</sub> = H, alkyl, tigloyl, benzoyl, or any other organic ester group;

R<sub>2</sub> = H or alkyl;

R<sub>3</sub> = H or OH;

In which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;

and in which the dashed lines indicate the optional presence of a further bond in the C4-C5 or C5-C6 positions;

[0029] The invention will be illustrated by the following Examples:

#### Example 1

[0030] The ability of the compound 3-O-[-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one-hereinafter referred to as P57AS3 - to prevent aspirin-induced damage to the stomach was investigated in the rat. Three groups each containing eight rats were given aspirin in 0.5% CMC orally at a dose of about 600 mg/kg (1.0 ml of 60 mg/ml). The first group (the control group) was additionally given 0.5% CMC. The second group was given compound PS7AS3 at a dose rate of 100mg/kg; and the third group (comparative group) was given cimetidine at a dose rate of 20mg/kg.

[0031] After 5 hours, the animals' stomachs were visually inspected for damage. Damage was assessed on a scale in which each small haemorrhage scored one point; each large haemorrhage scored two points; each small ulcer scored three points; and each large ulcer scored four points. The results obtained were as follows:

Table 1

Oral treatment	Dose (mg/kg)	Rat no.	Gastric damage score
Vehicle (0.5% CMC)	-	1	10
		2	10
		3	16
		4	8
		5	5
		6	3
		7	1
		8	6
		Mean	7.38
P57AS3	100	9	0
		10	0
		11	3
		12	0
		13	0
		14	0
		15	2
		16	0
		Mean	0.63
Cimetidine	20	17	0
		18	0
		19	4
		20	0
		21	0
		22	0
		23	0
		24	0
		Mean	0.50

[0032] These results show that compound P57AS3 gave an average reduction in gastric damage of 91.5% compared to 93.2% for cimetidine. In both cases, the statistical significance of the difference from the control group was  $P < 0.01$ .

#### Example 2

[0033] The effect of both the spray dried extract (sap from *Hoodia gordonii*) - referred to hereinafter as P57SD - and the purified compound P57AS3 were examined for effects on gastric acid secretion in the pylorusligated rat. Male Sprague-Dawley derived CD rats, 175-230 g, were used. On the day before the test, rats were outfitted with stainless steel tail cups to prevent coprophagy, housed individually with water allowed ad lib, and fasted overnight (18 hr). In one test the spray dried sap P57SD (5 or 50 mg/kg), or cimetidine (10 mg/kg) or vehicle (water) was orally administered to the rats immediately before fasting. Water was removed 30 minutes prior to ligation on the morning of the test. Under isoflurane anesthesia, a midline celiotomy was performed, and the pylorus of each rat was ligated with 4-0 silk.



[0034] In a second experiment, purified compound P57AS3 was administered intraduodenally (i.d.) at 1 or 10 mg/kg at the time of the pylorus ligation. In this experiment with intraduodenal administration of P57AS3, vehicle (DMSO), cimetidine (10 mg/kg) or P57AS3 (1 or 10 mg/kg) was administered immediately after ligation. The incisions were then closed with staples. In the experiments with subcutaneous administration of P57, vehicle (DMSO), cimetidine (10 mg/kg) or P57, 10 or 30 mg/kg were injected subcutaneously in the nape of the neck immediately after the incisions were closed. Two hours after ligation, the rats were killed, the contents of each stomach collected and cleaned by centrifugation (30 minutes at 2000 x g), and the volume of gastric fluid in each stomach was determined. Using a Radiometer pH meter and autotitrator, the pH and acid concentration of each gastric sample was determined by titration to pH 7 with 0.1N NaOH. Results are expressed as  $\mu\text{eq/hr/100 g}$  (body weight) and % vehicle control.

[0035] In the first experiment, gastric acid output was determined 18 hours after rats were orally administered the spray dried sap P57SD. Results showed that the 5 mg/kg dose of P57SD was without effect, but the 50 mg/kg dose inhibited gastric acid secretion by 40% (Table 2). In a second experiment, the purified compound P57AS3 was administered intraduodenally (i.d.) at 1 or 10 mg/kg at the time of the pylorus ligation. The 10 mg/kg i.d. dose of P57AS3 decreased acid secretion by 88%, but the 1 mg/kg i.d. dose had no effect (Table 2). Experiments were then conducted to determine if systemically administered P57AS3 would also inhibit gastric acid secretion. P57AS3 was administered subcutaneously at the time of ligation in 3 experiments using 10 mg/kg and in one experiment using 30 mg/kg. In one of the experiments with 10 mg/kg, P57AS3 produced a significant decrease in gastric acid secretion, but in the other 2 experiments with 10 mg/kg and the one experiment with 30 mg/kg there was a trend toward decreased acid output but the amount of acid secretion was not significantly different from vehicle-treated controls (Table 2). In each pylorus ligation experiment, gastric acid secretion was significantly reduced (range 36% to 83% inhibition) in a group of rats treated with cimetidine (10 mg/kg), which was included as a positive control.

Table 2

Effect of P57 on gastric acid secretion in the pylorus-ligated rat				
Exp.	Active	Dose (mg/kg)	Route	Acid Output % Control mean $\pm$ sd
I	P57SD	5	po	85.4 $\pm$ 34.5
		50	po	*60.0 $\pm$ 21.7
II	P57AS3	1	id	109.0 $\pm$ 37.8
		10	id	*12.4 $\pm$ 1.1
III	P57AS3	10	sc	48.9 $\pm$ 29.2
IV	P57AS3	10	sc	74.0 $\pm$ 38.0
V	P57AS3	10	sc	*57.1 $\pm$ 19.8
VI	P57AS3	30	sc	63.7 $\pm$ 17.3

\*significantly different from vehicle group,  $p \leq 0.05$

## SUMMARY OF RESULTS

### In Vitro Studies

[0036] Purified P57AS3 was examined for binding affinity to a variety of receptors, channels, and uptake sites in vitro. At a concentration of

10  $\mu\text{M}$ , in isolated tissue studies, P57AS3 at concentrations  $\leq 10 \mu\text{M}$  had no effect on the rate of spontaneously beating guinea pig right atria or on the force of contraction of electrically stimulated left atria. At concentrations of 1  $\mu\text{M}$  and higher, P57AS3 produced intermittent and transient elevations in the basal tension of electrically stimulated left atria. P57AS3 had no effect on resting tension of isolated guinea pig gallbladder, ileum or trachea, or on CCK octapeptide-induced contractions of the gallbladder (CCK-A receptor mediated), histamine induced contractions of the guinea pig ileum ( $H_1$  receptor mediated), or acetylcholine-induced contractions of the guinea pig ileum ( $M_3$  receptor mediated). At concentrations  $\geq 1 \mu\text{M}$ , P57AS3 noncompetitively inhibited carbachol-induced contractions of the guinea pig trachea ( $M_3$  receptor mediated).

## In Vivo Studies

[0037] The spray dried form of P57 was examined for effects on gastrointestinal motility and both the spray dried and purified forms were examined for effects on gastric acid secretion in the rat. At a dose of 5 mg/kg p.o., spray dried P57 had no effect on gastrointestinal transit or on gastric acid secretion. However, a 50 mg/kg dose of spray dried P57 inhibited gastric emptying by 26% and acid output by 40%. Following intraduodenal (id) administration in the rat, purified P57AS3 had no effect on gastric acid secretion at a dose of 1 mg/kg, but a 10 mg/kg dose decreased acid output by 88%. In 2 of 3 experiments at a dose of 10 mg/kg and in one experiment at a dose of 30 mg/kg, subcutaneously (sc) administered purified P57AS3 had no effect on gastric acid output, but in 1 of 3 experiments the 10 mg/kg sc dose reduced gastric acid secretion by 43%.

## References


[0038]

1. Hill SJ, and Young JM. Characterization of [3H]mepyramine binding to the longitudinal muscle of guinea pig small intestine. *Mol Pharmacol* 1981; 19:379-387
2. Eltze M, Mutschler E, and Lambrecht G. Affinity profiles of pizotifen, ketotifen and other tricyclic antimuscarinics at muscarinic receptor subtypes M1, M2 and M3. *Eur J Pharmacol* 1992; 211:283-293.
3. Bishop LA, Gerskowitch VP, Hull RA, Shankley NP, Black JW. Combined dose-ratio analysis of cholecystokinin receptor antagonists, devazepide, lorglumide and loxiglumide in the guinea pig gallbladder. *Br J Pharmacol* 1992; 106:61-66.

## Claims

1. Use of:

- (a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or
- (b) a compound of the formula (I)



wherein:

- R = an alkyl group containing from one to four carbon atoms;  
 R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;  
 R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

in which the acyl group R-(C=O)- group can be in the reduced form R-(C-OH)-;  
 and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position;  
 in the manufacture of a medicament for preventing or reducing a disorder of the alimentary system in an animal.

2. The use according to claim 1, wherein said medicament is for preventing or reducing a condition or disorder caused by or exacerbated by gastric acid secretion in an animal.
3. The use according to claim 1 or 2, wherein said disorder is selected from: excess gastric acid secretion, reflux oesophagitis, epigastric pain, dyspepsia, gastric or duodenal ulceration, acid aspiration, gastroduodenal erosion.
4. The use according to any preceding claim, wherein said disorder is gastric damage associated with use of a non-steroidal anti-inflammatory drug.
5. The use according to claim 4, wherein the medicament protects the gastro-intestinal tract of a mammal from damage caused by the non-steroidal anti-inflammatory drug.
6. The use according to any preceding claim, wherein in the general formula (I), said organic ester group is tigloyl, benzoyl or anthraniloyl.
7. The use according to any preceding claim, wherein in the general formula (I), R is an alkyl group having from one to four carbon atoms, R<sub>1</sub> is tigloyl or anthraniloyl; and a double bond is present in the C5-C6 position.
8. The use according to any preceding claim, wherein in the general formula (I), R<sub>2</sub> is a trisaccharide group.
9. The use according to claim 9, wherein the component sugars of said trisaccharide group are 6-deoxy and/or 2,6-dideoxy hexoses.
10. The use according to claim 9, wherein the terminal hexose moiety is thevetosyl.
11. The use according to any preceding claim, wherein the compound is 3-O-[-β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a derivative thereof.
12. The use according to any preceding claim, wherein said extract comprises spray-dried sap of the plant *Hoodia gordonii*.
13. The use according to any preceding claim, wherein said animal or said mammal is a human.
14. A pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a compound of the formula (I) as defined in any of claims 1 and 6 to 12 above.
15. A pharmaceutical composition as claimed in claim 14, which is prepared in unit dosage form.



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 00 30 9271

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X, D	WO 98 46243 A (ABRAMS MICHAEL JOHN ; CSIR (ZA); MAHARAJ VINESH (ZA); VLEGGAAR ROBE) 22 October 1998 (1998-10-22) * page 1, line 1 - page 4, line 28 *	1-15	A61K35/78 C07J7/00 C07J17/00
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			A61K
The present search report has been drawn up for all claims			
Place of search	Date of completion of the search	Examiner	
THE HAGUE	21 March 2001	Rempp, G	
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone  Y : particularly relevant if combined with another document of the same category  A : technological background  O : non-written disclosure  P : intermediate document</p> <p>T : theory or principle underlying the invention  E : earlier patent document, but published on, or after the filing date  D : document cited in the application  L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 30 9271

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

21-03-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9846243     A	22-10-1998	AU     7061398 A	11-11-1998
		BG     103795 A	31-05-2000
		BR     9808593 A	23-05-2000
		CN     1252000 T	03-05-2000
		EP     0973534 A	26-01-2000
		GB     2338235 A	15-12-1999
		HU     0000838 A	28-10-2000
		JP     2000510482 T	15-08-2000
		NO     994992 A	14-12-1999
		PL     336498 A	03-07-2000
		TR     9902540 T	21-01-2000
<hr/>			

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

